

注意：

允許學生個人、非營利性的圖書館或公立學校合理使用本基金會網站所提供之各項試題及其解答。可直接下載而不須申請。

重版、系統地複製或大量重製這些資料的任何部分，必須獲得財團法人臺北市九章數學教育基金會的授權許可。

申請此項授權請電郵 ccmp@seed.net.tw

Notice:

Individual students, nonprofit libraries, or schools are permitted to make fair use of the papers and its solutions. Republication, systematic copying, or multiple reproduction of any part of this material is permitted only under license from the Chiuchang Mathematics Foundation.

Requests for such permission should be made by e-mailing Mr. Wen-Hsien SUN ccmp@seed.net.tw

**International Mathematics
TOURNAMENT OF THE TOWNS**

Junior A-Level Paper

Fall 2008.

1. On a 100×100 chessboard, 100 Queens are placed so that no two attack each other. Prove that if the board is divided into four 50×50 subboards, then there is at least one Queen in each subboard.
2. Each of four stones weighs an integral number of grams. Available for use is a balance which shows the difference of the weights between the objects in the left pan and those in the right pan. Is it possible to determine the weight of each stone by using this balance four times, if it may make a mistake of 1 gram either way in at most one weighing?
3. Serge has drawn triangle ABC and one of its medians AD . When informed of the ratio $\frac{AD}{AC}$, Elias is able to prove that $\angle CAB$ is obtuse and $\angle BAD$ is acute.
Determine the ratio $\frac{AD}{AC}$ and justify your result.
4. Baron Münchhausen asserts that he has a map of Oz showing five towns and ten roads, each road connecting exactly two cities. A road may intersect at most one other road once. The four roads connected to each town are alternately red and yellow. Can this assertion be true?
5. Let $a_1, a_2, \dots, a_n$ be positive numbers such that $a_1 + a_2 + \dots + a_n \leq \frac{1}{2}$. Prove that $(1 + a_1)(1 + a_2) \cdots (1 + a_n) < 2$.
6. ABC is a non-isosceles triangle. E and F are points outside triangle ABC such that $\angle ECA = \angle EAC = \angle FAB = \angle FBA = \theta$. The line through A perpendicular to EF intersects the perpendicular bisector of BC at D . Determine $\angle BDC$.
7. In the infinite sequence $\{a_n\}$, $a_0 = 0$. For $n \geq 1$, if the greatest odd divisor of n is congruent modulo 4 to 1, then $a_n = a_{n-1} + 1$, but if the greatest odd divisor of n is congruent modulo 4 to 3, then $a_n = a_{n-1} - 1$. The initial terms are 0, 1, 2, 1, 2, 3, 2, 1, 2, 3, 4, 3, 2, 3, 2 and 1.
 - (a) Prove that the number 1 appears infinitely many times in this sequence.
 - (b) Prove that every positive integer appears infinitely many times in this sequence.

Note: The problems are worth 4, 6, 6, 6, 8, 9 and 5+5 points respectively.